


DNA-TR-82-179-V14

NUCLEAR WEAPON EFFECT RESEARCH AT PSR-1982

Volume XIV—Acute Radiation Response in Humans: Informal
Comments by Physicians and Radiobiologists

George H. Anno
Pacific-Sierra Research Corporation
12340 Santa Monica Boulevard
Los Angeles, California 90025



1 June 1983

Technical Report

CONTRACT No. DNA 001-82-C-0046

Distribution limited to US Government Agencies only;
Administrative/Operational Use, 13 February 1984. Other
requests for this document must be referred to Director,
Defense Nuclear Agency, Washington, DC 20305.

THIS WORK WAS SPONSORED BY THE DEFENSE NUCLEAR AGENCY
UNDER RDT&E RMSS CODE B384082466 V99QAXNL00097 H2590D.

Prepared for

Director

DEFENSE NUCLEAR AGENCY

Washington, DC 20305

DTL-84-2067

Destroy this report when it is no longer needed. Do not return to sender.

PLEASE NOTIFY THE DEFENSE NUCLEAR AGENCY,
ATTN: STTI, WASHINGTON, D.C. 20305, IF
YOUR ADDRESS IS INCORRECT, IF YOU WISH TO
BE DELETED FROM THE DISTRIBUTION LIST, OR
IF THE ADDRESSEE IS NO LONGER EMPLOYED BY
YOUR ORGANIZATION.



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER DNA-TR-82-179-V14	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) NUCLEAR WEAPON EFFECT RESEARCH AT PSR-1982 Volume XIV—Acute Radiation Response in Humans: Informal Comments by Physicians and Radiobiolo- gists		5. TYPE OF REPORT & PERIOD COVERED Technical Report
7. AUTHOR(s) George H. Anno		6. PERFORMING ORG. REPORT NUMBER PSR Report 1317
9. PERFORMING ORGANIZATION NAME AND ADDRESS Pacific-Sierra Research Corporation 12340 Santa Monica Boulevard Los Angeles, California 90025		8. CONTRACT OR GRANT NUMBER(s) DNA 001-82-C-0046
11. CONTROLLING OFFICE NAME AND ADDRESS Director Defense Nuclear Agency Washington, DC 20305		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Task V99QAXNL-00097
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE 1 June 1983
		13. NUMBER OF PAGES 32
		15. SECURITY CLASS (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE N/A since UNCLASSIFIED
16. DISTRIBUTION STATEMENT (of this Report) Distribution limited to US Government Agencies only; Administrative/ Operational Use, 13 February 1984. Other requests for this document must be referred to Director, Defense Nuclear Agency, Washington, DC 20305.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES This work was sponsored by the Defense Nuclear Agency under RDT&E RMSS Code B384082466 V99QAXNL00097 H2590D.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Ionizing Radiation Effects Acute Symptomatology Human Performance		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This report summarizes informal comments made by individuals of the medical/ radiobiological community regarding the response of humans to acute ionizing radiation effects. Experience with radiation therapy patients, nuclear accident victims, and animal experimentation are discussed in terms of acute radiation symptoms, physical and mental behavior, and the ability of individ- uals to perform military tasks if exposed to nuclear weapon radiation.		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

PREFACE

This report is one of a 16 volume set comprising the Pacific-Sierra Research Corporation (PSR) final report on Defense Nuclear Agency contract DNA 001-82-C-0046. The work done under this contract spans a wide range of nuclear weapon effect research covering airblast, cratering and ground motion, low-dose radiation, underground test design and development, fire research, and electromagnetic pulse research. The contract technical monitor was Cyrus P. Knowles.

This volume continues PSR's investigation of nuclear radiation effects on military troop performance for the Defense Nuclear Agency.* It presents information gathered in discussions with physicians and radiotherapists to complement and update prior work. This task was supervised by David L. Auton.

*Reported in G. H. Anno, H. L. Brode, and R. Washton-Brown, *Initial Human Response to Nuclear Radiation*, Defense Nuclear Agency, Washington, D.C., DNA-TR-81-237, 1 April 1982.

TABLE OF CONTENTS

PREFACE	1
 Section	 Page
Clarence E. Lushbaugh	3
Staff of Fred Hutchinson Cancer Research Center, Division of Oncology	5
Karel Dicke and Axel Zander	8
Robert E. George	9
George W. Casarett	10
Louis A. Gottschalk	13
Karl F. Hubner	14
Herbert Gerstner	15
John E. Pickering	15
Thomas Strike	17
Charles Turbyfill	18
Victor Bond	18
Alfred Bruner	19
Eugene Saenger	20
REFERENCES	21

ACUTE RADIATION RESPONSE IN HUMANS: INFORMAL COMMENTS
BY PHYSICIANS AND RADIOBIOLOGISTS

Pacific-Sierra Research Corporation is investigating the effects of intermediate radiation doses (100 to 3000 rads, free-in-air) on troop combat effectiveness. In the first phase of this research, we reviewed the literature and analyzed relevant data to develop models of symptomatic response as a function of dose, postexposure time, and symptom severity [Anno et al., 1982]. Before proceeding with the next research phase--to estimate how symptoms impair the physical and mental tasks associated with combat--we visited various specialists and research centers to obtain current data. Our informants included authors of the literature, radiation therapist physicians, and radiobiologists. This note summarizes the substance of their remarks. We have made no attempt to relate the summaries to each other and have arranged them in no particular order.

The visits, which took place from August to November 1981, considerably enriched our information base, and we intend to continue them in the future.

CLARENCE E. LUSHBAUGH, MEDICAL SERVICES DIVISION,
OAK RIDGE ASSOCIATED UNIVERSITIES,
OAK RIDGE, TENNESSEE

Fifty rads is the minimum dose at which noticeable symptoms^{*} of radiation sickness can be expected. At that level, however, sickness is detectable only by cytologic examination; there are no outward manifestations. At 100 rads, prodromal effects are visible in persons who are hypersensitive to radiation.[†] Therapy patients irradiated with

^{*}Throughout this note, "symptoms" is used to mean both subjective evidence and objective signs of radiation sickness. All doses are mid-line values unless specified otherwise.

[†]The categories hypersensitive, normosensitive, and hyposensitive are defined in Anno et al. [1982].

doses of at least 150 rads become intolerant of exercise or more easily fatigable, even in the absence of gastrointestinal symptoms. At 200 rads, most exposed persons manifest fever and other prodromal symptoms. At 300 rads (bone marrow dose), hemopoietic depression can be severe; blood counts must be closely watched so that

- Platelets do not fall below $20,000/\text{mm}^3$.
- White blood cells do not fall below $1000/\text{mm}^3$.
- Lymphocytes do not fall below 200 to $500/\text{mm}^3$.

All persons exposed to 300 rads experience nausea and vomiting, and many develop a painful hemorrhagic sore throat, a syndrome known as agranulocytic angina, which stems from pharyngeal ulceration, bacterial invasion, granulocytopenia, and thrombocytopenia. Doses above 400 rads exceed the LD_{50} level, and victims normally require hospitalization and medical care to survive. Persons subjected to 500 rads definitely require hospitalization and medical care to survive; a 600 rad dose may be the $LD_{50/60}$ level, even with hospital and medical care.

At 600 rads, the heart and blood vessels begin to be affected. At 1000 to 2000 rads, degeneration of the vascular endothelium allows blood fluids to leak outside vessel walls, blistering the skin. At 6000 rads, sclerotic processes cause irreparable vascular damage and gangrene sets in. Even at 1000 rads, only one in a million victims could survive without hospitalization and medical care such as fluids and bone marrow transplants; with hospital care, a few more could be saved.

At 1000 rads, most victims would suffer severe prodromal symptoms that would impair their performance of combat tasks. The lowering of blood pressure (hypotension) due to shock would further reduce performance. However, at that level, incapacitation might not occur until several hours postexposure, as with the accident victim exposed to 1114 rads [Hemplemann, Lisco, and Hoffman, 1952].

Blood vessel damage leading to cardiovascular shock syndrome is perhaps the most lethal and irreversible threat. Loss of 14 or 15 liters of serum and electrolyte fluids through leakage into extravascular tissues causes extreme circulatory problems, severe edema,

extracranial pressure, and cerebral anoxia, which can bring death within 2 days. Deaths attributed to central nervous system (CNS) failure at doses of less than 8000 rads are really caused by cardiovascular shock, whose symptoms resemble those of CNS failure. True CNS death, due to neurocytotoxic damage, does not occur below doses of 8000 rads.

Nausea, vomiting, and fatigability could affect performance. Nausea is much more debilitating than vomiting. The pyloric spasms of vomiting relieve stomach pressure, although too much vomiting, of course, disturbs electrolyte balance. Fatigability can be a serious problem, but we know very little about how it arises from exposure to radiation. It does not seem to develop immediately after exposure.

STAFF OF FRED HUTCHINSON CANCER RESEARCH CENTER,
DIVISION OF ONCOLOGY,
SEATTLE, WASHINGTON

Because of the differences between hospital and battlefield environments, it is a problem to extrapolate from the radiation responses of therapy patients to those of healthy young soldiers in combat. Before irradiation, therapy patients at this center are given 120 mg of cyclophosphamides to reduce the number of leukemic cells. The medication causes a lot of vomiting and some gastrointestinal distress; radiation treatment is delayed until symptoms have subsided somewhat, about 3 to 4 days. We do not know how such premedication affects radiation response--perhaps it aggravates gastrointestinal symptoms.

Therapy patients also differ from troops exposed to radiation in that all but a few are given fractional doses of 200 to 225 rads/day over 6 to 7 days. Total dosages amount to 1200 to 1575 rads, the latter given to leukemia patients in relapse. Dose rates are low, 5 to 8 rads/min. Extrapolation to the high dose rates and single exposures expected on the battlefield requires adjustments for the effects of different dose rates. Such adjustments are feasible if nausea, vomiting, and other symptoms have the same relationship of dose rate to response manifested in gastrointestinal-response and cell-survival experiments.

From experiments in which dogs given bone marrow transplants died of acute toxicity, the following dose-rate effects have been established:

<u>Dose Rate</u> <u>(rads/min)</u>	<u>Dose (rads)</u>
2.5	1800
5.0	1400
10.0	1000
20.0	800-1000

The dose-versus-dose-rate curve flattens out as predicted by cell-survival experiments, in which steep gradients were seen between 1 and 10 rads/min. At high dose rates, the dose-rate effect rapidly becomes less significant.

Prior chemotherapy does not seem to affect initial nausea and vomiting; 1 to 1.5 hr after receiving doses of 500 to 600 rads, most patients vomit, whether they have had chemotherapy or not. Nausea and intermittent vomiting per se may not severely affect performance, if the responses of irradiated monkeys hold true for humans. However, since any movement aggravates nausea and vomiting, continuation of combat operations after exposure might make those symptoms severe enough to impair battlefield performance. For therapy patients, of course, movement is minimized.

Within a few hours of irradiation, most patients show swelling of the parotid (parotitis), which lasts 24 to 48 hr. This effect, closely resembling mumps, can be quite painful.

High spiking temperatures, 40° to 40.5°C (104° to 104.9°F), are seen particularly in leukemia patients within a few hours of exposure and generally subside after 12 to 14 hr. The fever is not due to infection but is a response to massive cell breakdown, primarily of bone marrow cells. Such temperature elevations would have a debilitating effect on normal, healthy humans.

At high doses, the metabolic breakdown of cells, proteins, and amino acids can cause massive releases of uric acid. Without the administration of intravenous fluids to maintain the proper pH level,

crystals can form in the kidney tubules and cause irreversible renal shutdown 24 to 48 hr after radiation exposure. In another 24 to 48 hr, creatinine and potassium levels will rise and death will occur from cardiac arrest and other complications. An account of a case in which the foregoing sequence occurred appears in Thomas et al. [1971].

It is difficult to infer fatigability effects in exposed troops from those in irradiated leukemia patients because medical treatments such as chemotherapy, drugs to counter graft-versus-host disease (GVHD), and bone marrow transplants may affect fatigability.

About 10 percent of therapy patients contract a liver-related disease 6 to 21 days after irradiation. It is characterized by occlusion of the central venules, portal hypertension, and massive ascites resulting in pulmonary edema, fluid retention, and usually severe electrolyte fluid imbalance. Even with massive medical management it is difficult to get fluids recirculating, and the mortality rate is fairly high. Renal shutdown can also result.

Observations suggest that man and monkeys compare reasonably well in some responses to radiation. Soon after receiving a dose of 950 rads given at 7 rads/min (total radiation time, 2.25 hr), both begin vomiting; both recover after about 24 hr. Monkeys seem much more active after exposure, but the activity may be due to adrenalin release rather than radiation response per se, since they show fear when approached. We do not know whether high spiking temperatures or parotitis also occur in monkeys. The lung problems and pneumonitis seen in humans several months after radiation are not seen in monkeys or dogs.

In therapy patients treated with single doses of up to 1200 rads and fractionated doses of up to 1600 rads given over 7 days, no cardiovascular syndrome [Fanger and Lushbaugh, 1967] has been noticed. Nor has that syndrome been manifested in dogs exposed to single doses of 1600 to 1800 rads. Therapy patients have not shown hypotension, although the dose rates were perhaps not high enough to cause it (5 to 8 rads/min); hypotension occurred in monkeys after doses of 800 to 1000 rads given at ~200 rads/min [Chapman and Young, 1968].

For troops on the battlefield, doses of 1000 rads or perhaps as low as 600 rads would produce serious incapacitation within an hour or two that would last 48 hr. Without intensive medical care, fluids, and antiemetics, problems brought on by severe fluid loss, tissue damage, and elevated temperatures--such as electrolyte imbalance, glucose perturbation, uric acid buildup, and edema--would lead to rapid degeneration and death shortly thereafter.

KAREL DICKE AND AXEL ZANDER,
RADIATION THERAPY DEPARTMENT,
M. D. ANDERSON HOSPITAL AND TUMOR INSTITUTE,
HOUSTON, TEXAS

Leukemia patients who received single doses of 750 to 950 rads (with chemotherapy) had toxic reactions that threatened their survival. About 30 percent of those patients experienced severe pulmonary distress known as white lungs 7 to 10 days after radiation. It resulted in hypoxia, acute respiratory problems, and sometimes further complications ending in death, even with the use of a respirator. A significant increase in toxicity was observed when doses were increased from 750-800 rads to 900-950 rads. Toxicity problems also occurred in patients with relapsed leukemia who received doses of 1200 rads administered over a period of 3 days, in six fractions of 200 rads each. Problems included severe mucositis, fever, parotitis, pneumonia, infections, and other complications. All patients had received autologous bone marrow transplants.

Current therapy protocol requires six fractions of 170 rads each (given at 25 rads/min, i.e., within 6.8 min) for a total of 1020 rads. Patients in relapse and remission are exposed twice a day (a.m. and p.m., 6 hr apart) over a period of 3 days. All patients receive bone marrow transplants 24 to 48 hr after irradiation.

All patients experience diarrhea, which may be related to the chemotherapy and antibiotics they receive; 90 percent contract mucositis in varying degrees, most severely those receiving the anti-GVHD drug methotrexate; and 90 percent show the spiking fever. Patient survivability is good, much better than with the large single-dose regimen

mentioned above. The marked reduction in pneumonitis and lung distress may owe partially to the plastic isolation bubble, which allows patients to recover from treatment in an aseptic environment.

As for the time sequence of prodromal symptoms, all patients become nauseated and vomit 0.5 to 1 hr after the first radiation fraction. After the second fraction, symptoms are more severe than after the first, but severity lessens increasingly after the third and subsequent fractions. There is some diarrhea following the fifth and sixth fractions, according to nurses in the ward.

A 29-year-old male patient in remission verified the foregoing prodromal response sequence. This patient also exercised daily throughout his treatment. Besides experiencing nausea, vomiting, and stomatitis, he became quite fatigued while exercising the first day; the fatigability continued for several days, diminishing somewhat a week or so after the last fraction.

If we could extrapolate directly from clinical experience to the battlefield, and assume that performance impairment depends on symptom severity, we would expect that a soldier could not drive a tank after exposure to 340-680 rads (corresponding to the cumulative therapeutic dose received the second day) but could drive it after exposure to 680-1020 rads (corresponding to the third and fourth days). However, such direct extrapolation would be fallacious because it may only be the medical care found in a hospital that enables exposed persons to respond well after absorbing larger amounts of radiation.

ROBERT E. GEORGE,
DEPARTMENT OF RADIATION ONCOLOGY,
INDIANA UNIVERSITY HOSPITALS,
INDIANAPOLIS, INDIANA

Most of the treatment in this therapy department is with half-body irradiation at midline doses of 600 to 700 rads. The main purpose is to control pain in patients whose cancer has spread to the bone from the prostate or breast. Responses to half-body irradiation are of course less severe than those to total-body irradiation (TBI). And it

is difficult to isolate the degree of debilitation resulting from radiation per se since the patients are quite sick to begin with. However, parotid swelling, nausea, and vomiting are usually seen in all irradiated patients.

Work at the Armed Forces Radiobiological Research Institute (AFRRI)* tested the early transient incapacitation (ETI) effects of radiation on animal neurological systems. In separate experiments, monkeys and miniature pigs were trained to perform simple tasks (such as crossing a shuttle box on signal), then tested in those tasks immediately after irradiation. Both experiments showed that the animals were less incapacitated after exposure to neutron than to gamma radiation at high doses and dose rates. Those surprising results suggest that performance is impaired by a mechanism other than just cell killing, at which neutrons are known to be more effective. Accordingly, the neutron relative biological effectiveness (RBE) may be less than unity for considerations of performance impairment though greater than unity for lethality.

After receiving a 1000 rad dose, a soldier might still be able to point or fire a gun, at least for a short while; however, he would tire quickly and his endurance would decline markedly. More complicated tasks--like aiming and firing a missile system or flying an airplane--might be a problem soon after exposure.

GEORGE W. CASARETT,
RADIATION BIOLOGY AND BIOPHYSICS DEPARTMENT,
UNIVERSITY OF ROCHESTER MEDICAL CENTER,
ROCHESTER, NEW YORK

The RBE for neutrons is based on comparison with a low linear energy transfer (LET) standard radiation (e.g., X-rays or gamma rays) for a particular biological reaction. RBE is also a function of dose, increasing as dose decreases. For fast neutrons in the moderate to high dose range, the RBE is between 3 and 5 for many biological end

* Before assuming his present position, Dr. George was chairman of the radiation biology department at AFRRI.

points; it would be more like 3 for early (prodromal) responses. In the low to very low dose range, the RBE for neutrons may be much higher. Nearly all data on systemic radiation effects in various tissues show that neutrons create much more damage than gamma radiation. It is well known that high LET radiation such as fast neutrons is much more lethal for entire animal organisms as well as more effective at killing cells.

Some nonradiation insults might be used to simulate radiation damage and study its effects on physical performance. Candidates for such experimentation would be insults that waste muscle tissue, cause it to "give up" rapidly, or change the creatinine/creatine ratio. For muscle performance, radiation probably affects endurance much more than short-term exertion.

"Pure cardiovascular" death from radiation, as described by Fanger and Lushbaugh [1967], may not exist. Cardiovascular effects always accompany the gastrointestinal syndrome beginning around 500 R (and resulting in death at about 1000 R) and the CNS syndrome beginning at about 2000 R (with death at about 5000 R). The degree of cardiovascular damage of course depends on the dose. A complicating factor in isolating cardiovascular effects is that the gastrointestinal syndrome, which may occur 5 to 14 days after exposure, overlaps with hemopoietic depression. Even the hemopoietic syndrome ending in death always has some associated cardiovascular problems.

Experiments with rats indicate that the key to gastrointestinal syndrome severity is the amount of radiation exposure to the gut. In some animals the gut was surgically moved to the exterior, irradiated, then replaced in the nonirradiated abdomen; in other animals the entire abdomen, with gut in place, was irradiated. For all, death occurred only 1 day apart; after 5 days for animals with the reimplanted guts, and after 4 days for the others.

As for the CNS response, the rat experiments showed that large doses of several thousand rads delivered only to the head produced essentially the same acute CNS symptoms leading to incapacitation and death as did equivalent whole-body doses.

Acute responses to TBI in man may be classified as in Table 1 (adapted from Rubin and Casarett [1968]).

Table 1. Acute effects of TBI in humans.

Item	Syndrome		
	Central Nervous System	Gastrointestinal	Hemopoietic
Organ most affected	Brain	Small intestine	Bone marrow
Syndrome threshold dose	2000 R	500 R	100 R
Postexposure time of onset	1/4-3 hr	3-5 days	2-3 weeks
Death threshold dose	5000 R	1000 R	200 R
Postexposure time of death	Within 2 days	3-14 days	3 weeks-2 months
Characteristic symptoms	Lethargy Tremors Convulsions Ataxia	Malaise Anorexia Nausea Vomiting Diarrhea Gastrointestinal malfunction Fever Dehydration Electrolyte loss Circulatory collapse	Malaise Fever Dyspnea on exertion Fatigue Leukopenia Thrombopenia Purpura
Major underlying pathology	Vasculitis Encephalitis Meningitis Edema	Depletion of intestinal epithelium Neutropenia (marrow damage) Infection	Bone marrow atrophy Pancytopenia Infection Hemorrhage Anemia

LOUIS A. GOTTSCHALK,
DEPARTMENT OF PSYCHIATRY AND HUMAN BEHAVIOR,
COLLEGE OF MEDICINE, UNIVERSITY OF CALIFORNIA,
IRVINE, CALIFORNIA

Speech content analysis was used to study the effect of total-body and half-body irradiation on human cognitive and emotional processes [Gottschalk et al., 1969]. Sixteen patients with metastatic carcinoma were exposed to doses of 50 to 300 R cobalt 60 radiation as palliative treatment. After exposure the patients were encouraged to talk on any subject that came to mind, and 5 min samples of their remarks were tape-recorded.

Most simply, content analysis involves determining the frequency, in a sample of speech, of words judged to be of research interest. Gottschalk and his colleague G. C. Gleser use the clause as the primary unit of analysis and emphasize the context of the sample. The analysis focuses not only on the structure (syntax) of the clause and the meaning (semantics) of individual words and phrases but also on the emotional tone of the clause as a whole.

Technicians "scored" the samples by detecting references to concepts and actions reflecting the patient's emotional state. References could take many forms, from direct citation of a certain word to metaphorical allusion in several clauses. Accurate and consistent scoring required careful and time-consuming training, even for technicians well versed in the language. Computerized techniques were developed to automate the procedure.

According to an intellectual impairment scale developed by Gottschalk and Gleser, the nine patients for whom data were complete showed evidence of transiently impaired intellectual function ($P \leq 0.02$). The impairment appeared immediately after irradiation and lasted one day. But the difference between impairment after actual radiation and after sham radiation was statistically insignificant, which raises the question whether that effect was only a result of irradiation. Criteria other than the researchers' impairment scale showed no worsening of mentation after either actual or sham radiation.

In the entire patient group, transient anxiety was significantly greater before treatment (actual or sham) than afterward ($P \leq 0.02$).

The data also provide objective evidence that hopefulness is higher in patients who have been hospitalized a shorter time ($P \leq 0.01$) and have a longer expected survival time ($P \leq 0.08$).

KARL F. HUBNER, MEDICAL SERVICES DIVISION,
OAK RIDGE ASSOCIATED UNIVERSITIES,
OAK RIDGE, TENNESSEE

The radiation responses of accident victims and therapy patients are tempered by the hospitalization and intensive medical care they receive after irradiation. In extrapolating those responses to the battlefield, clinically observed effects should be expected at lower doses. How much lower is a matter of uncertainty. Part of the uncertainty might be resolved by careful animal experimentation, but a scaling relationship would have to be developed between humans and animals, perhaps one based on lifespan. An anchor point is the $LD_{50/60}$ level of ~325 rads, postulated by Lushbaugh, which reflects no medical care. With medical care, that level would be reached at substantially higher doses, perhaps 500 to 600 rads.

It is not clear what battlefield medical care could be provided to counter damage to the hemopoietic system. No experimental work has evaluated the effect of whole-blood transfusions. Granulocytes might not be available, although isolation in a sterile environment would help reduce infection. Intravenous fluids would be essential.

Marshall Brucer, formerly chief of the medical division of the Oak Ridge National Laboratory, has developed three-dimensional response-model concepts (response versus dose and time) for a variety of observable acute effects, including LD_{50} , vomiting, elevated body temperature, lowered white blood cell and platelet counts, fatigue, and psychological upset [Brucer, comp., 1959].

The National Cancer Institute, American Cancer Society, and other medical organizations have recommended scales for gauging the extent of illness based on performance. Such a scale, like the Karnofsky scale, might be useful in relating radiation sickness to performance.

HERBERT GERSTNER, OAK RIDGE, TENNESSEE

Dr. Gerstner, now retired, was formerly with the Air Force School of Aerospace Medicine. His remarks focused on studies of radiation responses in therapy patients at M. D. Anderson Hospital and Tumor Institute in Houston. The studies were based on a detailed reconstruction of nurses' records over a 2-year period and a dosimetry evaluation by Warren Sinclair.

Patients exposed to 100 rad TBI twice a week became nauseated and vomited about 30 min after treatment, but those symptoms disappeared within 24 hr. Patients then remained asymptomatic for another 3 weeks, after which their white blood cell count fell.

Animal experiments as well as the M. D. Anderson studies suggest that it is possible to roughly classify an exposed population by the severity of members' symptoms and their sensitivity to radiation. Hypersensitive patients will show symptoms after doses as low as 50 rads; everyone will be affected after 400 rads. Only 1 out of 99 would have a chance of surviving after receiving 500 rads. At a dose of 100 to 150 rads, about 20 percent of the population will be unaffected, 20 percent mildly affected, 50 percent moderately affected, and 10 percent severely affected. As the dose increases, the distribution will shift toward greater severity. About 20 percent of an exposed population will be hyposensitive to radiation, 20 percent hypersensitive, and 60 percent normosensitive or average.

The severity of symptoms or sensitivity to radiation exhibited in the initial (prodromal) phase is not necessarily proportional to severity or sensitivity exhibited in the secondary (manifest-illness) phase; completely different mechanisms are operating.

JOHN E. PICKERING,
U.S. AIR FORCE SCHOOL OF AEROSPACE MEDICINE,
BROOKS AIR FORCE BASE, TEXAS

Researchers here are investigating how well an aircrew might be able to perform the various phases of a flight mission (e.g., takeoff, climbout, cruise, refueling) after exposure to nuclear radiation. To

define those abilities, they seek predictions of performance decrement with respect to dose and time after exposure along a mission-time profile [Albanese and Pickering, 1974]. Gastrointestinal distress is likely to be the first acute radiation sickness symptom to accompany, if not cause, performance decrement.

The opinions of persons with experience as rated crew members and those familiar with nuclear weapon radiation effects have yielded useful judgments about the likelihood of an aircrew's accomplishing each mission phase.* And the case histories of two accident victims provide data for gauging the effects of supralethal doses (2000 to 5000 rads). However, the most realistic data come from flight simulation tests using trained, irradiated monkeys. Monkeys display essentially the same clinical responses to radiation as humans: vomiting, fatigue, lethargy, erythema, and epilation.

Adolescent rhesus monkeys were trained in flight-simulation tasks involving control-precision dynamics (pitch and roll) and visual, auditory, and memory responses. In one experiment, their performance of the tasks was measured immediately after and several hours after irradiation with 600 rads of GODIVA neutrons. The radiation was delivered in pulses of 50 to 100 μ secs nearly uniformly over the whole body. The estimates of an aircrew's postirradiation vulnerability noted below are based on analysis of the results of that experiment. A second experiment exposing monkeys to 1050-1100 rads is underway.

During the first 4 hr after exposure, the monkeys presented a clinical picture like that observed in the human accident victims studied. The animals were lethargic, actually lying on the floor of their cage, were nonaggressive, nonresponsive to food, and suffered episodes of vomiting. Nevertheless, they performed their tasks on the day of exposure and succeeding days for the required 4 hr, with alternating 12 min work periods and 3 min rest periods. The number of

*If differences of opinion diminish when experts are polled repeatedly, there is greater probability that the converging judgments are accurate [Dalkey, 1969].

performance errors increased in 5 of the 8 subjects on exposure day and 2 of the 8 subjects on each test day thereafter. Reaction times increased in 7 of the 8 subjects on exposure day, in 3 of the 8 subjects on the second day, 5 of the 8 subjects on the third day, and 4 of the 8 subjects on the fourth day.

Extrapolating to human beings, the researchers judge that a flight crew exposed to 600-650 rads could probably complete a strategic B-52 mission in the expected 12 to 24 hr period because some recovery will take place during that time.

In light of the incident with the carrier *Nimitz*, it might be a problem for a crew to complete a Navy light attack mission requiring a carrier landing within 4 hr of a nuclear attack. It might be prudent to bring the aircraft near the carrier and have the crew "punch out." That procedure would be clearly indicated if the crew had been exposed to more than 600 rads.

F-14 and F-16 crews, encumbered by flight gear and chemical protective garments, would probably be unable to make a second tactical sortie if exposed to more than 400 rads.

The most stringent limitation applies to the Airborne Command Post. Its crew may be required to complete a flight in a fallout field within 4 hr of a nuclear attack, land somewhere, then "regenerate" 4 hr later. With that mission, and to ensure totally alert, decisive, symptom-free personnel, crew members should have been exposed to no more than 100 rads/hr (yielding a cumulative dose of 200 rads in 12 hr and 237 rads in 24 hr).

THOMAS STRIKE,
NATIONAL INSTITUTES OF HEALTH,
BETHESDA, MARYLAND

It has been suggested that postirradiation impairment of cognitive performance results from a drop in blood pressure triggered by release of antihistamines in the blood. Experiments were carried out at the AFRRRI to determine the effect of antihistamine medication on performance impairment after radiation exposure. The medication significantly

reduced performance impairment as well as hypotension. Researchers at Walter Reed Army Medical Center are developing antiradiation drugs, the most common of which is WR 2721.

CHARLES TURBYFILL,
NATIONAL INSTITUTES OF HEALTH,
BETHESDA, MARYLAND

Monkeys are reasonable human surrogates for radiation response testing; monkey data fit human data fairly well. After doses of 10,000 rads and more, monkeys die in about 7 to 8 hr. Performance is impaired 1.6 to 2 min after exposure and for 6 to 7 min thereafter, followed by 3 to 4 hr of apparently normal ability. Then, performance impairment resumes and continues until death. After doses of 400 to 600 rads, performance impairment is fairly small.

The antiradiation drug developed by M. H. Hieffer and D. E. Davidson, Jr. at Walter Reed Army Medical Center protects the sulfa-hydral (SH) group against radiation damage. It must be administered before irradiation.

There was an account of a Japanese general at Hiroshima who force-marched his troops away from the city after the bombing; the account should be found and studied for information on military performance after radiation exposure.

VICTOR BOND,
BROOKHAVEN NATIONAL LABORATORY,
LONG ISLAND, NEW YORK

In studying the acute effects of radiation exposure on performance, it is important to identify precisely what tasks are being tested and how performance is measured--even if only binary measurement is possible (i.e., subject can/cannot perform X task). Measurements of muscle strength and endurance can be made by monitoring bicycle exercise with an ergometer. Sports medicine specialists are familiar with the various kinds of physical performance measurements.

Useful data on postirradiation performance could be obtained from staff nurses caring for patients being treated with radiation therapy or suffering from other toxic conditions such as uremia or alcohol poisoning. Continual close contact with patients enables nurses to make keen judgments about the abilities of patients affected by various symptoms.

Acute radiation effects seem to depend on a dose threshold; exposure above the threshold leads to radically different behavior than does exposure below the threshold. If performance ability can be correlated with the nature and intensity of specific symptoms, similar correlations could be expected with performance impairment.

ALFRED BRUNER,
NATIONAL INSTITUTES OF HEALTH,
BETHESDA, MARYLAND

Ten years of extensive experimentation with monkeys^{*} suggest the following conclusions regarding human radiation responses. Within a few minutes of exposure to as little as 300 rads of cobalt 60 radiation (or the equivalent) at a dose rate of 30 rads/min (whole body), personnel may be expected to undergo transient performance decrement/incapacitation if the task is a complex one. Higher dose rates will increase the incidence and severity of the decrement; simpler tasks will reveal less impairment. Cardiovascular homeostatic mechanisms are compromised briefly after sufficient exposure, the most noticeable effect being a peripheral relaxation-hypotensive shock syndrome. A subsequently developing cerebral hypoxia is the presumed basis for performance decrement/incapacitation, which arises from a complex composite of constitutional, situational, and radiation parameters not entirely clear. Any additional external demands imposed on the cardiovascular system at this time, such as exercise or high G-force flight maneuvers, may be expected to further jeopardize brain oxygenation and therefore

^{*}Conducted while Dr. Bruner was at the Lovelace Foundation for Medical Education and Research, Albuquerque, New Mexico. Experimental results are summarized in Bruner [1977a and 1977b].

performance integrity. The histamine hypothesis of early postirradiation effects suggests that protection against the syndrome may be afforded by preirradiation blocking of H_1 and H_2 histamine receptors through drug pretreatment. It might be useful to evaluate simulated postirradiation performance in human volunteers by administering drugs that depress cardiovascular functioning and produce nausea during task performance.

EUGENE SAENGER,
UNIVERSITY OF CINCINNATI MEDICAL CENTER,
CINCINNATI, OHIO

Improvements in medical care over the last decade have produced a rapidly expanding data base from radiation therapy that can be exploited for studying the effects of radiation exposure. It may be inappropriate, however, to apply radiation response models obtained from TBI therapy patients to the battlefield. Troops will probably receive partial- not total-body irradiation, and the dose distribution in their bodies will probably vary widely from soldier to soldier.

Early performance may not be severely impaired at doses of 600 rads and less. Performance impairment will probably occur early and be severe at doses of 1200 rads and more. It is the performance effects of intermediate doses--600 to 1200 rads--that are not well understood.

REFERENCES

- Albanese, R. A., and J. E. Pickering, "Aircrew Vulnerability in Nuclear Encounters," *Mil. Med.*, Vol. 13, No. 12, 1974, pp. 945-951.
- Anno, G. H., H. L. Brode, and R. Washton-Brown, *Initial Human Response to Nuclear Radiation*, Defense Nuclear Agency, Washington, D.C., DNA-TR-81-237, 1 April 1982 (also published by Pacific-Sierra Research Corporation, Note 477, April 1982).
- Brucer, M. B. (comp.), *The Acute Radiation Syndrome: A Medical Report on the Y-12 Accident, June 16, 1958*, U.S. Atomic Energy Commission, Report ORINS-25, April 1959.
- Bruner, A., "Immediate Changes in Estimated Cardiac Output and Vascular Resistance after ^{60}Co Exposure in Monkeys: Implications for Performance Decrement," *Radiat. Res.*, Vol. 70, 1977, pp. 391-405 (1977a).
- , "Immediate Dose-Rate Effects of ^{60}Co on Performance and Blood Pressure in Monkeys," *Radiat. Res.*, Vol. 70, 1977, pp. 378-390 (1977b).
- Chapman, P. H., and R. Young, "Effect of Cobalt-60 Gamma Irradiation on Blood Pressure and Cerebral Blood Flow in the Macaca Mulatta," *Radiat. Res.*, Vol. 35, 1968, pp. 78-85.
- Dalkey, N. C., *The Delphi Method: An Experimental Study of Group Opinion*, The Rand Corporation, Santa Monica, California, RM-5888-PR, 1969.
- Fanger, H., and C. C. Lushbaugh, "Radiation Death from Cardiovascular Shock Following a Criticality Accident," *Arch. Path.*, Vol. 83, May 1967, pp. 446-460.
- Gottschalk, L. A., et al., "Total and Half Body Irradiation, Effect on Cognitive and Emotional Processes," *Arch. Gen. Psychiat.*, Vol. 21, November 1969, pp. 574-580.
- Hemplemann, L. H., H. Lisco, and J. G. Hoffman, "The Acute Radiation Syndrome: A Study of Nine Cases and a Review of the Problems," *Ann. Int. Med.*, Vol. 36, 1952, pp. 279-570.
- Rubin, P., and G. W. Casarett, *Clinical Radiation Pathology*, W. B. Saunders Company, Philadelphia, 1968.
- Thomas, E. D., et al., "Allogeneic Marrow Grafting for Hematologic Malignancy Using HL-A Matched Donor-Recipient Sibling Pairs," *Blood*, Vol. 38, No. 3, September 1971, pp. 267-287.

DISTRIBUTION LIST

DEPARTMENT OF DEFENSE

Armed Forces Staff College
ATTN: Library

Assistant Secretary of Defense
International Security Affairs
ATTN: ISA/PP
ATTN: F. Miller
ATTN: Policy Plans & NSC Affairs

Assistant Secretary of Defense
Policy Analysis
ATTN: D. Shilling

Assistant to the Secretary of Defense
Atomic Energy
ATTN: Executive Assistant
ATTN: R. Wagner

Commander-in-Chief, Atlantic
ATTN: J22
ATTN: J3

Commander-in-Chief, Pacific
ATTN: J-54
ATTN: C3SRD

Defense Advanced Rsch Proj Agency
ATTN: TTO
ATTN: T. Bache

Defense Communications Agency
ATTN: Code J300, M. Scher

Defense Intelligence Agency
ATTN: DB 4C2, D. Spohn
ATTN: DT, Sci-Tech Intell
ATTN: Library
ATTN: DB
ATTN: DB-4C
ATTN: DN
ATTN: DB-4N
ATTN: DT, J. Vorona
ATTN: DB-1, Rsch, Sov Wpn Div, G. Ferrell
ATTN: RTS-2C, Tech Svcs & Spt
ATTN: DIO-GPF, W. Magathan
ATTN: DE, Estimates

Defense Nuclear Agency
ATTN: RAAE
ATTN: NAFD
ATTN: STSP
ATTN: NATA
ATTN: RAEV
ATTN: STRA
ATTN: SPSS
ATTN: SPTD
ATTN: STNA
ATTN: SPSS, G. Ullrich
ATTN: NATD
ATTN: RAEV
ATTN: STBE
ATTN: K. Schwartz
ATTN: NASD
4 cy ATTN: STTI/CA

DEPARTMENT OF DEFENSE

Defense Technical Information Center
2 cy ATTN: DD

Deputy Under Sec of Def, S&TNF
ATTN: T. Jones

Deputy Under Sec of Def Policy
ATTN: R. Stivers

Field Command
DNA, Det 1
Lawrence Livermore National Lab
ATTN: FC-1

Field Command
DNA, Det 2
Los Alamos National Lab/DST
ATTN: MS-635 FC-2

DNA PACOM Liaison Office
ATTN: J. Bartlett

Field Command
Defense Nuclear Agency
ATTN: FCPR
ATTN: FCPRK, R. Wells
ATTN: FCTT
ATTN: FCTT, W. Summa
ATTN: FCTXE
ATTN: FCTK, C. Keller

Interservice Nuclear Weapons School
ATTN: TTV
ATTN: Document Control

Joint Chiefs of Staff
ATTN: J-5 Strategy Div, W. McClain
ATTN: SAGA/SSD
ATTN: SAGA/SFD
ATTN: J-5 Nuc/Chem Policy Br, J. Steckler
ATTN: J-3 Strategic Operations Division
ATTN: J-5 Nuclear Division/Strategy Div

Joint Data System Support Ctr
ATTN: C-312, R. Mason

Joint Strat Tgt Planning Staff
ATTN: JPTM
ATTN: JLK, DNA Rep
ATTN: JPST
ATTN: JP
ATTN: JLKS
ATTN: JLKC

National Defense University
ATTN: NWCLB-CR

Office of the Sec of Defense
Net Assessments
ATTN: Document Control

Principal Dep Under Sec of Defense
Research & Engineering
ATTN: J. Wade, Jr

DEPARTMENT OF DEFENSE (Continued)

Program Analysis & Evaluation
ATTN: Strategic Programs
ATTN: S. Johnson

US European Command
ATTN: ECJ-3

US European Command
ATTN: ECJ-5

US National Military Representative
SHAPE

Attention US Doc Ofc for
ATTN: Intel
ATTN: OPS, Nuc Concepts
ATTN: Nuc Plans

US Readiness Command
ATTN: J-3

Under Sec of Defense for Policy
ATTN: Dir Negotiations Policy, S. Buckley
ATTN: F. Ikle
ATTN: Dir Strategic Policy, C. Estes
ATTN: Dir Ping & Requirements, M. Sheridan

Under Secy of Def for Rsch & Engrg
ATTN: R. Delauer
ATTN: Strat & Space Sys (OS)
ATTN: Strat & Arms Control, L. Menichiello
ATTN: Strat & Theater Nuc for B. Stephan
ATTN: Strat & Space Sys (OS), C. Knowles
ATTN: K. Hinman

United States Central Command
ATTN: CCJE-03, Daigneault

DEPARTMENT OF THE ARMY

Asst Ch of Staff for Intelligence
ATTN: DAMI-FIT

BMD Advanced Technology Center
ATTN: 1CRDABH-X
ATTN: ATC-T

Chief of Engineers
Department of the Army
ATTN: DAEN-MPE-T
ATTN: DAEN-RDL

Dep Ch of Staff for Ops & Plans
ATTN: DAMO-NCN
ATTN: DAMO-SSM, Pol-Mil Div
ATTN: Technical Advisor
ATTN: DAMO-RQS
ATTN: DAMO-RQA, Firepower Div
5 cy ATTN: DAMO-NC, Nuc Chem Dir

Harry Diamond Laboratories
ATTN: DELHD-NP
ATTN: DELHD-TA-L
ATTN: DELHD-DE
ATTN: 00100 Commander/Tech Dir/Div Dir
ATTN: DELHD-TD, Tech Dir
ATTN: Chief Div 30000
ATTN: DELHD-NW-P

DEPARTMENT OF THE ARMY (Continued)

US Army Armament Rsch Dev & Cnd
ATTN: DRDAR-LCN-E

US Army Ballistic Research Labs
ATTN: DRDAR-BLV
ATTN: DRDAR-BLA-S
ATTN: R. Reisler
ATTN: DRDAR-BLT, J. Keefer
ATTN: DRDAR-BLV-R, Dr Rainis

US Army Chemical School
ATTN: ATZN-CM-CC

US Army Comb Arms Combat Dev Acty
ATTN: ATZL-CAP

US Army Comd & General Staff College
ATTN: DTAC
3 cy ATTN: Combined Arms Research Library

US Army Concepts Analysis Agency
ATTN: CSSA-ADL

US Army Engineer Ctr & Ft Belvoir
ATTN: DT-LRC

US Army Engineer Div Huntsville
ATTN: HNEED-SR

US Army Engineer Div Ohio River
ATTN: ORDAS-L

US Army Engineer School
ATTN: Library

US Army Engr Waterways Exper Station
ATTN: J. Zelasko
ATTN: J. Strange
ATTN: J. Houston
ATTN: WESSE
ATTN: Library
ATTN: WESSA, W. Flathau
2 cy ATTN: WESSD, J. Jackson

US Army Foreign Science & Tech Ctr
ATTN: DRXST-SD-1

US Army Infantry Ctr & Sch
ATTN: ATSH-CD-CSO

US Army Intel Threat Analysis Det
ATTN: IAX-ADT

US Army Intelligence Center & School
ATTN: ATSI-CD-CS

US Army Logistics Center
ATTN: ATCL-OSS, S. Cockrell

US Army Material & Mechanics Rsch Ctr
ATTN: Technical Library

US Army Material Dev & Readiness Cnd
ATTN: DRCDE-D
ATTN: DRXAM-TL

DEPARTMENT OF THE ARMY (Continued)

US Army Materiel Sys Analysis Actvy
ATTN: XS, W3JCAA

US Army Mobility Equip R&D Cmd
ATTN: DRDME-WC, Technical Lib (Vault)

US Army Nuclear & Chemical Agency
ATTN: MONA-OPS
ATTN: Library
ATTN: MONA-OPS, B. Thomas
ATTN: MONA-OPS, J. Ratway

US Army TRADOC Sys Analysis Actvy
ATTN: ATAA-TAC

US Army Training and Doctrine Comd
ATTN: ATCD-FA

US Army War College
ATTN: AMCAC, F. Braden, Dept of Tactics
ATTN: Library
ATTN: War Gaming Facility

USA Military Academy
ATTN: Document Library

USA Missile Command
ATTN: DRSMI-XF
ATTN: Document Section
ATTN: DRSMI-RH

DEPARTMENT OF THE NAVY

Anti-Submarine Warfare Sys Proj Ofc
ATTN: PM-4

Charleston Naval Shipyard
ATTN: Commanding Officer

David Taylor Naval Ship R&D Ctr
ATTN: Code 142-3, Library
ATTN: Code 1750, J. Sykes
ATTN: Code 1750, W. Conley
ATTN: Code 174

Joint Cruise Missiles Project Ofc
ATTN: JCNG-707

Marine Corps
ATTN: DCS, (P&O) Requirements Div
ATTN: Code OTD0-31
ATTN: DCS (P&O) Strategic Plans Div

Marine Corps Dev & Education Command
ATTN: Commander

Naval Air Development Center
ATTN: Code 702, B. McHugh

Naval Air Systems Command
ATTN: Code 350D, H. Benefiel

Naval Civil Engineering Laboratory
ATTN: Code L-51, S. Johnson
ATTN: Code L-51, J. Crawford

Naval Electronic Systems Command
ATTN: PME 117-21

DEPARTMENT OF THE NAVY (Continued)

Naval Facilities Engineering Command
ATTN: Code 04B

Naval Intelligence Command
ATTN: NIC-01

Naval Intelligence Support Ctr
ATTN: NISC-40
ATTN: NISC-30

Naval Material Command
ATTN: MAT 08T-22
ATTN: MAT-00
ATTN: MAT-046

Naval Ocean Systems Center
ATTN: R. Hammond
ATTN: J. Hooper

Naval Postgraduate School
ATTN: Code 56PR
ATTN: Code 1424 Library
ATTN: G. Lindsay

Naval Research Laboratory
ATTN: Code 2627

Naval Sea Systems Command
ATTN: SEA-06H2
ATTN: SEA-406
2 cy ATTN: SEA-6431G, H. Seguire

Naval Submarine School
ATTN: Commanding Officer

Naval Surface Weapons Center
ATTN: Code R44, H. Glaz
ATTN: Code R14
ATTN: Code R41, N. Scofield
ATTN: Code U41
ATTN: Code F31
ATTN: Code F30

Naval Surface Weapons Center
ATTN: Code DG-502, E. Freiling
ATTN: Tech Library & Info Svcs Br

Naval War College
ATTN: Code E-11, Tech Service

Naval Weapons Center
ATTN: Code 32607, L. Thompson

Naval Weapons Evaluation Facility
ATTN: H. Struve
ATTN: J. Abbott
ATTN: Code 10
ATTN: Technical Director
ATTN: G. Binns

Navy Field Operational Intelligence Office
ATTN: Commanding Officer

Newport Laboratory
ATTN: K. Walsh
ATTN: Code EM, J. Kalinowski

DEPARTMENT OF THE NAVY (Continued)

Nuclear Weapons Tng Group, Atlantic
ATTN: Nuclear Warfare Department

Nuclear Weapons Tng Group, Pacific
ATTN: Nuclear Warfare Department

Ofc of the Deputy Chief of Naval Ops
ATTN: NOP 985F
ATTN: NOP 951, ASW Div
ATTN: NOP 955, AAW Div
ATTN: NOP 021
ATTN: NOP 022
ATTN: NOP 06
ATTN: NOP 09
ATTN: NOP 950, Force Level Plns Div
ATTN: NOP 953, Tac Readiness Div
ATTN: NOP 954, Strike & Amphib Warf Div
ATTN: NOP 32, Surf Warf Div
ATTN: NOP 35, Surf Cbt Sys Div
ATTN: NOP 50, Avn Plns & Rqmts Dev
ATTN: NOP 963
ATTN: NOP 654, Strat Eval & Anal Br
ATTN: NOP 05
ATTN: NOP 987
ATTN: NOP 03
ATTN: NOP 02
ATTN: NOP 098, Ofc Res-Dev-Test & Eval
ATTN: NOP 981
ATTN: NOP 03EG
3 cy ATTN: NOP 96, N Prog Ofc-Sys Anal Div

Office of Naval Research
ATTN: Code 474, N. Perrone
ATTN: Code 431
ATTN: Code 200

CNO Executive Panel
Office of the Chief of Naval Operations
ATTN: OP-00K

Strategic Systems Project Office
ATTN: NSP-43

Surface Warfare Development Group
ATTN: Commander

Surface Warfare Officers School Cmd
ATTN: Combat Systems Dept

DEPARTMENT OF THE AIR FORCE

Air Force
ATTN: INE, Estimates

Air Force
ATTN: INT

Air Force Geophysics Laboratory
ATTN: LMH, H. Ossing

Air Force Institute of Technology
ATTN: Library
ATTN: ENP

Air Force Office of Scientific Rsch
ATTN: J. Allen
ATTN: W. Best

DEPARTMENT OF THE AIR FORCE (Continued)

Air Force Operational Test & Eval Ctr
ATTN: OA
ATTN: OAY, Capt Lutz

Air Force Propulsion Lab
ATTN: LKDH, Stop 24, E. Haberman

Air Force Systems Command
ATTN: DLW

Air Force Weapons Laboratory
ATTN: NT
ATTN: NTE, M. Plamondon
ATTN: NTES, J. Thomas
ATTN: NTES-C, R. Henny
ATTN: SUL
ATTN: NXS
ATTN: NTYC, J. Burgio

Air University Library
ATTN: AUL-LSE

Assistant Chief of Staff
Studies & Analysis
2 cy ATTN: AF/SAMI, Tech Info Div

Ballistic Missile Office/DAA
ATTN: ENSN, A. Schenker
ATTN: ENSN, E. Furbee
ATTN: ENMP, D. Van Gari
ATTN: PP
ATTN: ENMP
ATTN: EN
2 cy ATTN: ENSN

Deputy Chief of Staff
Research, Development & Acq
ATTN: AFRDQA
ATTN: AFRDQI
4 cy ATTN: AFRD-M, Spec Asst for MX

Deputy Chief of Staff
Plans and Operations
ATTN: AFXOOR, Opns, Opnl Spt
ATTN: Dir of Plans, AFXOX
ATTN: AFXOXFM, Plns, Frc Dev Mun Plns

Foreign Technology Division
ATTN: SD
ATTN: NIIS Library
ATTN: TQ

Pacific Air Forces
ATTN: IN
ATTN: XO

Rome Air Development Center
ATTN: TSLD

Strategic Air Command
ATTN: XPO
ATTN: XPFS
ATTN: INAO
ATTN: NRI/STINFO Library

US Air Force Scientific Advisory Bd
ATTN: AF/NB

DEPARTMENT OF THE AIR FORCE (Continued)

Tactical Air Command
ATTN: TAC/DR
ATTN: TAC/INO

DEPARTMENT OF ENERGY

Albuquerque Operations Office
ATTN: R. Jones
ATTN: CTID

Office of Military Application
ATTN: OMA, DP-22

Nevada Operations Office
ATTN: Doc Con for Technical Library

Division of Reactor Rsch and Dev
ATTN: P. Hemmig

OTHER GOVERNMENT AGENCIES

Central Intelligence Agency
ATTN: Office of Soviet Affairs
ATTN: Office of East Asia
ATTN: Natl Intel Council/Anal Gp, H. Ford
ATTN: Office of Global Issues
ATTN: OSR/SE/F
ATTN: OSMR/NED

Department of Commerce
National Bureau of Standards
ATTN: J. Hubell

Department of the Interior
Bureau of Mines
ATTN: Tech Lib

Federal Emergency Management Agency
ATTN: Ofc of Rsch/NP, D. Bensen
ATTN: Asst Assoc Dir for Rsch, J. Kerr

NORAD
ATTN: JSYS, F. Smith

Office of Technology Assessment
ATTN: M. Harris

US Arms Control & Disarmament Agcy
ATTN: A. Lieberman
ATTN: H. Cooper

DEPARTMENT OF ENERGY CONTRACTORS

University of California
Lawrence Livermore National Lab
ATTN: L-21, M. Gustavson
ATTN: Technical Info Dept Library
ATTN: L-71, R. Howerton
ATTN: L-8, F. Barrish
ATTN: D. Glenn
ATTN: L-389, R. Andrews
ATTN: L-531, A. O'Dell
ATTN: L-35, J. Immele
ATTN: R. Werne
ATTN: R. Corallo

DEPARTMENT OF ENERGY CONTRACTORS (Continued)

Los Alamos National Laboratory
ATTN: R. Sandoval
ATTN: J. Hopkins
ATTN: F. Begay
ATTN: R. Stolpe
ATTN: Reports Library
ATTN: MS 634, T. Dowler

Oak Ridge National Laboratory
Union Carbide Corp, Nuclear Div
ATTN: Civil Def Res Proj
ATTN: Rad Shielding Ctr
ATTN: Central Rsch Library
ATTN: C. Clifford

Sandia National Laboratories
ATTN: Lib & Sec Class Div

Sandia National Laboratories
ATTN: Org 7112, A. Chabal
ATTN: 0334, J. Struve
ATTN: Tech Lib 3141
ATTN: 0332, J. Keizur
ATTN: 0333, R. Stratton
ATTN: L. Hill

DEPARTMENT OF DEFENSE CONTRACTORS

Academy for Interscience Methodology
ATTN: N. Painter

Aerospace Corp
ATTN: Library
ATTN: Library Acquisition M1/199

Agbabian Associates
ATTN: M. Agbabian

Applied Research Associates, Inc
ATTN: J. Drake
ATTN: J. Bratton
ATTN: H. Auld
ATTN: M. Higgins

Applied Research Associates, Inc
ATTN: S. Blouin

Applied Research Associates, Inc
ATTN: D. Piepenburg

Applied Research Associates, Inc
ATTN: R. Frank

Applied Theory, Inc
2 cy ATTN: J. Trullo

AVCO Systems Division
ATTN: Library A830

BDM Corp
ATTN: F. Leech
ATTN: D. Percy

Boeing Co
ATTN: MS-85-20, D. Choate

DEPARTMENT OF DEFENSE CONTRACTORS (Continued)

BDM Corp

ATTN: L. Schlipper
ATTN: P. White
ATTN: R. Welander
ATTN: J. Braddock
ATTN: R. Buchanan
ATTN: C. Somers
ATTN: J. Bode
ATTN: J. Morgan
ATTN: J. Herzog
ATTN: Corporate Lib
ATTN: C. Masaff
ATTN: T. Neighbors
ATTN: H. Portnoy

Boeing Co

ATTN: A. Lowrey
ATTN: Aerospace Library

CACI, Inc - Federal

ATTN: A. Berry

California Institute of Technology

ATTN: T. Ahrens

California Research & Technology, Inc

ATTN: K. Kreyenhagen
ATTN: M. Rosenblatt
ATTN: S. Schuster
ATTN: Library

California Research & Technology, Inc

ATTN: D. Orphal
ATTN: F. Sauer

Calspan Corp

ATTN: Library

Carpenter Research Corp

ATTN: H. Carpenter

66th MI Group

ATTN: K. Moran

Data Memory Systems, Inc

ATTN: T. Dupuy

University of Denver

ATTN: Sec Officer for J. Wisotski

EG&G Wash Analytical Svcs Ctr, Inc

ATTN: Library

Electro-Mech Systems, Inc

ATTN: R. Shunk

GARD

ATTN: G. Heidhardt

Harold Rosenbaum Associates, Inc

ATTN: G. Weber
ATTN: H. Rosenbaum

Horizons Technology, Inc

ATTN: R. Kruger

Hudson Institute, Inc

ATTN: NAVMAG

DEPARTMENT OF DEFENSE CONTRACTORS (Continued)

IIT Research Institute

ATTN: M. Johnson
ATTN: Documents Library
ATTN: R. Welch

Inst for Foreign Anal, Inc

ATTN: J. Record

Inst for Foreign Pol Anal, Inc

ATTN: R. Faltzgraph

Institute for Defense Analyses

ATTN: Classified Library
ATTN: E. Kerlin
ATTN: Y. Utgoff

International Energy Associates, Inc

ATTN: L. Scheinman

IRT Corp

ATTN: Library

JAYCOR

ATTN: E. Almquist

Kaman Avidyne

ATTN: M. Hobbs
ATTN: Library

Kaman Sciences Corp

ATTN: Library

Kaman Sciences Corp

ATTN: E. Conrad

Kaman Sciences Corporation

ATTN: R. Miller

Kaman Tempo

ATTN: C. Anderson
ATTN: DASIAC

Kaman Tempo

ATTN: DASIAC

Lockheed Missiles & Space Co, Inc

ATTN: Technical Information Center
ATTN: T. Geers, Org 52-33

Lockheed Missiles & Space Co, Inc

ATTN: TIC-Library
ATTN: J. Weisner, Dept 80-82

Martin Marietta Denver Aerospace

ATTN: D-6074, G. Freyer

Maxwell Laboratories, Inc

ATTN: J. Murphy

McDonnell Douglas Corp

ATTN: Technical Library Services
ATTN: R. Halprin

McLean Research Center, Inc

ATTN: W. Schilling

University of Miami

ATTN: Contract Office, S. Wang

DEPARTMENT OF DEFENSE CONTRACTORS (Continued)

Merritt CASES, Inc
ATTN: J. Merritt
ATTN: Library

Natl Institute for Public Policy
ATTN: C. Gray

University of New Mexico
ATTN: M. Baum

ORI, Inc
ATTN: R. Wiles
ATTN: B. Buc

Orion Engineering, Inc
ATTN: W. Parks

Pacific-Sierra Research Corp
ATTN: G. Lang
ATTN: S. Finn
2 cy ATTN: H. Brode, Chairman SAGE
2 cy ATTN: B. Morgan, Jr
2 cy ATTN: L. Crusier
2 cy ATTN: D. Wilson
2 cy ATTN: G. Anno
2 cy ATTN: G. McClellan
2 cy ATTN: L. Schlessinger
5 cy ATTN: Library

Pacific-Sierra Research Corp
ATTN: D. Gormley
ATTN: G. Moe

Paulmar Corp
ATTN: C. Feldbau, B. Garrett, B. Glaser

Patel Enterprises, Inc
ATTN: M. Patel

Physics International Co
ATTN: L. Behrmann
ATTN: Technical Library
ATTN: E. Moore

R&D Associates
ATTN: J. Marcum
ATTN: F. Field
ATTN: W. Wright
ATTN: J. Lewis
ATTN: Tech Info Center
ATTN: R. Montgomery
ATTN: D. Simons
ATTN: P. Haas
ATTN: C. K. B. Lee
ATTN: A. Wohlstetter
ATTN: G. Jones
ATTN: G. Ivy

R&D Associates
ATTN: H. Polk
ATTN: A. Deverill
ATTN: J. Thompson

Radiation Research Associates, Inc
ATTN: Library

DEPARTMENT OF DEFENSE CONTRACTORS (Continued)

Rand Corp
ATTN: N. Levin
ATTN: J. Digby
ATTN: Library
ATTN: R. Solomon
ATTN: M. Pollack
ATTN: T. Parker
ATTN: P. Davis
ATTN: H. Gelman

Rand Corp
ATTN: B. Bennett
ATTN: T. Warner
ATTN: R. Goettmouller

S-CUBED
ATTN: T. Riney
ATTN: D. Grine
ATTN: Library

Science & Engineering Associates, Inc
ATTN: H. Linnerud

Science Applications, Inc
ATTN: M. Woolson
ATTN: J. Martin
ATTN: Technical Library
ATTN: M. Drake
ATTN: E. Straker

Science Applications, Inc
ATTN: T. Albert

Science Applications, Inc
ATTN: D. Maxwell
ATTN: D. Bernstein
ATTN: B. Dial

Science Applications, Inc
ATTN: W. Zimmerman
ATTN: P. Setty
ATTN: J. Goldstein
ATTN: W. Layson
ATTN: J. McGahan

Science Applications, Inc
ATTN: D. Kaul

Southwest Research Institute
ATTN: A. Wenzel
ATTN: M. Baker

SRI International
ATTN: G. Abrahamson
ATTN: D. Keough
ATTN: Y. Gupta
ATTN: J. Naar
ATTN: B. Lew
ATTN: W. Jaye

Structural Mechanics Associates, Inc
ATTN: R. Kennedy

SY Corp
ATTN: S. Weiss

DEPARTMENT OF DEFENSE CONTRACTORS (Continued)

System Planning Corp
ATTN: J. Luquier
ATTN: G. Parks

Systems Research & Applications Corp
ATTN: S. Greenstein

Teledyne Brown Engineering
ATTN: F. Leopard
ATTN: D. Ormond

Terra Tek, Inc
ATTN: J. Schatz
ATTN: A. Abou-Sayed
ATTN: S. Green
ATTN: Library

Tetra Tech, Inc
ATTN: L. Hwang

Tetra Tech, Inc
ATTN: F. Bothwell
ATTN: J. Preston

DEPARTMENT OF DEFENSE CONTRACTORS (Continued)

Titan Systems, Inc
ATTN: C. Albo

TRW Electronics & Defense Sector
ATTN: Technical Information Center
ATTN: P. Bhuta
2 cy ATTN: N. Lipner

TRW Electronics & Defense Sector
ATTN: P. Dai
ATTN: E. Wong

Universal Analytics, Inc
ATTN: E. Field

Weidlinger Assoc, Consulting Engrg
ATTN: M. Baron

Weidlinger Assoc, Consulting Engrg
ATTN: J. Isenberg

SUBJECT NAME FROM	Suspense DATE DATE	Number Type
	FILE DESIGNATION	

SUMMARY

4 up

TO	TO <i>AS</i>	TO	TO	TO
DATE <i>11/14/84</i>	DATE <i>11/15/84</i>	DATE	DATE	DATE
REPLIED OR INDORSED TO		FILED <i>(Place)</i>	OTHER ACTION <i>Peterson</i>	